



Last time

- More on Linear independence and spanning
 - Bases for subspaces
-

If $\{\vec{v}_1, \dots, \vec{v}_r\}$ is a basis for the subspace V , then any vector \vec{v} in V can be written uniquely as a linear combination

$$\vec{v} = c_1 \vec{v}_1 + \dots + c_r \vec{v}_r$$

Ex $\begin{bmatrix} x \\ y \end{bmatrix} = x \begin{bmatrix} 1 \\ 0 \end{bmatrix} + y \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ 

Ex $\begin{bmatrix} x \\ y \end{bmatrix} = x \begin{bmatrix} 1 \\ 0 \end{bmatrix} + \frac{y}{2} \begin{bmatrix} 0 \\ 2 \end{bmatrix}$ 

Ex $\begin{bmatrix} x \\ y \end{bmatrix} = x \begin{bmatrix} 1 \\ 1 \end{bmatrix} + (y-x) \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ 

As we saw last time,

columns of A linearly independent $\iff A\vec{x} = \vec{0}$ has a unique solution $\iff \ker A = \{\vec{0}\} \iff \text{rk}(A) = n$

columns of A span \mathbb{R}^m $\iff A\vec{x} = \vec{b}$ has a solution \vec{x} for every \vec{b} $\iff \text{im } A = \mathbb{R}^m \iff \text{rk}(A) = m$

Combining these, $\iff A$ is invertible

columns of A are a basis for \mathbb{R}^m $\iff A\vec{x} = \vec{b}$ has a unique solution for every \vec{b} $\iff \ker A = \{\vec{0}\}$ and $\text{im } A = \mathbb{R}^m \iff \begin{matrix} \text{rk}(A) \\ = \\ m \\ = \\ n \end{matrix}$

let's try to find a basis for the kernel and image of the matrix

$$A = \begin{bmatrix} 1 & 2 & 2 & -5 & 6 \\ -1 & -2 & -1 & 1 & -1 \\ 4 & 8 & 5 & -8 & 9 \\ 3 & 6 & 1 & 5 & 7 \end{bmatrix}$$

$$\text{RREF}(A) = \begin{bmatrix} 1 & 2 & 0 & 3 & -4 \\ 0 & 0 & 1 & -4 & 5 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

The general solution to $A\vec{x} = \vec{0}$ is

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = \begin{bmatrix} -2x_2 - 3x_4 + 4x_5 \\ x_2 \\ 4x_4 - 5x_5 \\ x_4 \\ x_5 \end{bmatrix}$$
$$= r \underbrace{\begin{bmatrix} -2 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}}_{\vec{v}_1} + s \underbrace{\begin{bmatrix} -1 \\ 0 \\ -5 \\ 0 \\ 0 \end{bmatrix}}_{\vec{v}_2} + t \underbrace{\begin{bmatrix} 4 \\ -6 \\ 5 \\ 0 \\ 4 \end{bmatrix}}_{\vec{v}_3},$$

So $\ker(A) = \text{Span}(\{\vec{v}_1, \vec{v}_2, \vec{v}_3\})$. But there can be no relations among these vectors, since each has a nonzero entry where the others don't.

$\implies \{\vec{v}_1, \vec{v}_2, \vec{v}_3\}$ is a basis for $\ker(A)$

In particular, $\dim \ker(A) = 3$.

The dimension of $\ker(A)$ is the number of free variables. Basis vectors are obtained by setting one free variable equal to 1 and all others to 0.

What about the image? We need to eliminate the redundant vectors from the columns of A . These correspond

to the redundant columns in $\text{RREF}(A)$,
even though these columns span
different subspaces.

$$A = \begin{bmatrix} 1 & 2 & 2 & -5 & 6 \\ -1 & -2 & -1 & 1 & -1 \\ 4 & 8 & 5 & -8 & 9 \\ 3 & 6 & 1 & 5 & 7 \end{bmatrix}$$

In an example, the
second, third and fourth
columns are clearly redundant, since e.g.,

$$\text{RREF}(A) = \begin{bmatrix} 1 & 2 & 0 & 3 & -4 \\ 0 & 0 & 1 & -4 & 5 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 3 \\ -4 \\ 0 \\ 0 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} - 4 \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}$$

Equivalently, the vector $\begin{bmatrix} -3 \\ 0 \\ 4 \\ 0 \end{bmatrix}$ is in

the kernel of $\text{RREF}(A)$. But this is the same as $\ker(A)$! So the same relation holds for the columns of A .

So a basis for the image is

$\left\{ \begin{bmatrix} 1 \\ -1 \\ 4 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ -1 \\ 5 \\ 1 \end{bmatrix} \right\}$ and $\dim \text{im}(A) = 2$.

The dimension of $\text{im}(A)$ is the rank of A . A basis is given by the columns of A that become pivot columns in $\text{rREF}(A)$