


Last time

- Sarrus' rule

- Patterns + inversions



- Determinants

$$\det(A) = \sum_P \text{sgn}(P) \text{prod}(P)$$

- Upper triangular case

Goal $\det(A) = 0 \iff A$ is not invertible

Idea We test invertibility with row

reduction, so let's try to relate determinants and row operations.

Based on the 2×2 case, we have a clear expectation.

Guess Let A be an $n \times n$ matrix.

(1) Swapping two rows of A changes the determinant by -1 .

(2) Scaling a row by k scales the determinant by k .

(3) Adding a multiple of a row to another doesn't change the determinant

Would this actually help us? If we can check this guess, then it will follow that

$\det(A) = 0 \iff \det(\text{RREF}(A)) = 0$,
since none of the row operations can
change whether $\det(A)$ is zero.

But $\text{RREF}(A)$ is upper triangular
with every diagonal entry either 0
or 1. So

$$\det(\text{RREF}(A)) = \begin{cases} 1 & \text{RREF}(A) = I_n \\ 0 & \text{otherwise.} \end{cases}$$

$$\det(A) \neq 0 \iff A \text{ is invertible}$$

Our guess will follow from two important properties of determinants.

(A) The determinant is "multilinear," i.e., linear in each row separately. Specifically, given fixed vectors $\vec{v}_1, \dots, \vec{v}_{i-1}, \vec{v}_{i+1}, \dots, \vec{v}_n \in \mathbb{R}^n$,



the function

$$T(\vec{x}) = \det \begin{bmatrix} \vec{v}_1 \\ \dots \\ \vec{v}_{i-1} \\ \vec{x} \\ \vec{v}_{i+1} \\ \dots \\ \vec{v}_n \end{bmatrix}$$

is linear. Indeed, for every pattern P , $\text{prod } P$ is a product of $(n-1)$ fixed numbers (drawn from the \vec{v} 's) with a single entry of \vec{x} . This is obviously linear in \vec{x} , so $\text{prod } P$ is, so $\det(A)$ is.

(B) The determinant is "alternating", i.e., $\det(A) = 0$ if two adjacent rows are equal.

Indeed, for every pattern P , there is another pattern P' that interchanges the entries drawn from the duplicate rows:



The entries of P' drawn from these rows are inverted precisely if they were not inverted in P , and all other inversions are the same (because the rows are adjacent). Hence $\text{prod}(P) = \text{prod}(P')$ and $\text{sgn}(P) = -\text{sgn}(P')$, so the terms cancel.

Back to row operations.

(1) Swapping two rows

$$\begin{aligned}
 0 &= \det \begin{bmatrix} \vdots \\ \vdots \\ x_u + x_v \dots \\ \vdots \\ x_v + x_u \dots \\ \vdots \end{bmatrix} = \det \begin{bmatrix} \vdots \\ \vdots \\ x_v \dots \\ \vdots \\ x_u \dots \\ \vdots \end{bmatrix} + \det \begin{bmatrix} \vdots \\ \vdots \\ x_u \dots \\ \vdots \\ x_v \dots \\ \vdots \end{bmatrix} \\
 &= \det \begin{bmatrix} \vdots \\ \vdots \\ x_v \dots \\ \vdots \\ x_u \dots \\ \vdots \end{bmatrix} + \det \begin{bmatrix} \vdots \\ \vdots \\ x_u \dots \\ \vdots \\ x_v \dots \\ \vdots \end{bmatrix} + \det \begin{bmatrix} \vdots \\ \vdots \\ x_u \dots \\ \vdots \\ x_v \dots \\ \vdots \end{bmatrix} + \det \begin{bmatrix} \vdots \\ \vdots \\ x_v \dots \\ \vdots \\ x_u \dots \\ \vdots \end{bmatrix}
 \end{aligned}$$

~~$$= \det \begin{bmatrix} \vdots \\ \vdots \\ x_u \dots \\ \vdots \\ x_v \dots \\ \vdots \end{bmatrix} + \det \begin{bmatrix} \vdots \\ \vdots \\ x_v \dots \\ \vdots \\ x_u \dots \\ \vdots \end{bmatrix}$$~~

If the rows aren't adjacent, swap adjacent rows until they are.

(2) Scaling a row

$$\det \begin{bmatrix} \vdots \\ \vdots \\ kx_i \\ \vdots \\ \vdots \end{bmatrix} = k \cdot \det \begin{bmatrix} \vdots \\ \vdots \\ x_i \\ \vdots \\ \vdots \end{bmatrix}$$

(3) Adding a multiple of a row

$$\det \begin{bmatrix} \vdots \\ \vdots \\ x_i + kx_j \\ \vdots \\ \vdots \end{bmatrix} = \det \begin{bmatrix} \vdots \\ \vdots \\ x_i \\ \vdots \\ \vdots \end{bmatrix} + k \det \begin{bmatrix} \vdots \\ \vdots \\ x_j \\ \vdots \\ \vdots \end{bmatrix} \rightarrow 0$$

As a final note, the determinant is also multilinear and alternating in the columns of the matrix.